

The Intersection of Two Fermat Hypersurfaces in $\mathbb{P}^3$ via Computation of Quotient Curves

Vijaykumar Singh* and Gary McGuire†

School of Mathematical Sciences

University College Dublin

Ireland

Email: vijaykumar.singh@ucdconnect.ie, gary.mcguire@ucd.ie

November 25, 2010

Mathematics Subject Classification: 11G20, 14H45

Abstract

We study the intersection of two particular Fermat hypersurfaces in $\mathbb{P}^3$ over a finite field. Using the Kani-Rosen decomposition we study arithmetic properties of this curve in terms of its quotients. Explicit computation of the quotients is done using a Gröbner basis algorithm. We also study the p -rank, zeta function, and number of rational points, of the modulo p reduction of the curve. We show that the Jacobian of the genus 2 quotient is $(4, 4)$ -split.

1 Introduction

Let $\mathbb{F}_q$ denote the finite field with q elements, where $q = p^n$ and p is a prime. The equations $x^a + y^a + z^a + w^a = 0$, $x^b + y^b + z^b + w^b = 0$ each define a Fermat hypersurface in $\mathbb{P}^3(\mathbb{F}_q)$, and their intersection defines a curve. This article studies some arithmetic properties of this curve in the particular case $a = 2$, $b = 3$, and $p \geq 5$. We therefore study the curve defined by the equations

$$x^2 + y^2 + z^2 + w^2 = 0, \quad x^3 + y^3 + z^3 + w^3 = 0,$$

over $\mathbb{Q}$ (and $\overline{\mathbb{Q}}$), by studying some of its quotient curves, and consider the reduction modulo p . It would be nice to have a formula for the number of rational points over $\mathbb{F}_p$, but our investigations show that such a formula is unlikely to exist.

*Research supported by Claude Shannon Institute, Science Foundation Ireland Grant 06/MI/006.

†Research supported by Claude Shannon Institute, Science Foundation Ireland Grant 06/MI/006.

Previous work on these curves has mostly been done when $p = 2$, because of connections to cyclic codes in algebraic coding theory. The case $a = 1$ and b arbitrary (when $p = 2$) was studied in [9]. The case $a = 1$ is somewhat different to the case $a > 1$ because when $a = 1$ the curve is a plane curve and its equation is easily obtained. Cases $a = 3, b = 5$ and $a = 1, b = 11$ are studied in [3], where the zeta function was computed and applied. The two above equations can be considered as a system of diagonal equations. Such systems have been studied before. Wolfmann [11] and Knapp [12] study a system with terms having the same exponents but different coefficients, over finite fields and local fields respectively.

In Section 2 we present some background needed for the paper. In Section 3 we give some relevant invariant theory, which is needed to provide the Kani-Rosen decomposition, and our algorithm to compute the Gröbner basis for a quotient curve, which may be new. Section 4 calculates the genus, proves nonsingularity, and computes the quotients involved in the Kani-Rosen decomposition. Section 5 studies the two genus 1 quotients, and Section 6 studies the genus 2 quotient. The genus 2 quotient exhibits some interesting behaviour, having a Jacobian which we prove is a $(4, 4)$ -split. These two sections work over $\overline{\mathbb{Q}}$. Section 7 considers the good reductions of our curve modulo primes, and the L-polynomials. Section 8 deals with the number of points on the curves over $\mathbb{F}_q$, and shows some data from computations.

Acknowledgements. We would like to thank Nils Bruin and Christophe Ritzenthaler for their helpful comments, and Alexey Zaytsev for many helpful discussions.

2 Background

In this section we present a summary of the background needed for this paper.

2.1 Algebraic Geometry

Let k be a field, $\mathbb{A}^n(k)$ be affine n -space. For $S \subset k[x_1, \dots, x_n]$ define

$$V(S) = \{x \in \mathbb{A}^n(k) \mid f(x) = 0 \quad \forall f \in S\}.$$

If S consists of a single (nonconstant) polynomial then $V(S)$ is called a hypersurface. A subset V of $\mathbb{A}^n(k)$ is called an affine algebraic set if $V = V(S)$ for some $S \subset k[x_1, \dots, x_n]$. Then all polynomials in $k[x_1, \dots, x_n]$ that vanish on V forms an ideal $I(V)$ called ideal of V . The affine coordinate ring $k[V]$ of V is quotient ring $k[x_1, \dots, x_n]/I(V)$. If V is irreducible or equivalently $I(V)$ is a prime ideal then V is called affine algebraic variety and the quotient of integral domain $k[x_1, \dots, x_n]/I(V)$ is called function field of V denoted by $K(V)$. If we allow only the homogenous polynomials in S we have a projective algebraic set and hence a projective variety. If for a variety C , the function field $k(C)$ has a transcendental degree 1 over k , then C is called an algebraic curve.

The genus of a curve is defined using the Riemann-Roch theorem. A non-singular curve of genus 1 is called an elliptic curve. A curve X of genus ≥ 2 is called hyperelliptic if there is a map $f : X \rightarrow \mathbb{P}^1$ of degree 2.

An algebraic group is a variety V together with a morphism $\oplus : V \times V \rightarrow V$ such that the set of points of V with the operation given by $\oplus$ is a group. A projective algebraic group is called an abelian variety.

Let A and B be two abelian varieties. The morphisms from A to B , which is also a group homomorphism forms a $\mathbb{Z}$ -module is denoted by $\text{Hom}(A, B)$. An element $\phi \in \text{Hom}(A, B)$ is called an *isogeny* if $\ker(\phi)$ is finite and $\text{Im}(\phi) = B$.

2.2 Curves

Let C be an algebraic curve defined over k . Recall that the *Jacobian variety* J_C is an abelian variety with the group structure corresponding the quotient group $\text{Pic}^0(C)$, the degree 0 divisors modulo the principal divisors.

The *zeta function* of a curve $C/\mathbb{F}_q$ is given by

$$Z(t, C) := \exp \left\{ \sum_{m=1}^{\infty} N_m \frac{t^m}{m} \right\}$$

where N_m is the number $\mathbb{F}_q^m$ -rational points. It was shown by Artin and Schmidt that $Z(t, C)$ can be written in the form

$$\frac{L(t)}{(1 - qT)(1 - T)}$$

where $L(t) \in \mathbb{Z}[t]$ (called the L -polynomial) is of degree $2g$, where g is the genus of the curve C . For any abelian variety $A = J_C$, the characteristic polynomial of the Frobenius endomorphism (acting on the l -adic Tate module of A), will be denoted $f_A(t)$. It is independent of l and has coefficients in $\mathbb{Z}$. In fact, Weil proved that $f_A(t)$ is the reciprocal polynomial of the L -polynomial of C .

We will use the following fundamental result.

Theorem 2.1. (Tate) *If A and B are the abelian varieties defined over $\mathbb{F}_q$. Then A is $\mathbb{F}_q$ -isogenous to abelian subvariety of B if and only if $f_A(t)$ divides $f_B(t)$ in $\mathbb{Q}[t]$. In particular, $f_A(t) = f_B(t)$ if and only if A and B are $\mathbb{F}_q$ -isogenous.*

Given an abelian variety A defined over k , the p -rank of A is defined by

$$r_p(A) = \dim_{\mathbb{F}_p} A(\bar{k})[p],$$

where $A(\bar{k})[p]$ is the subgroup of p -torsion points over the algebraic closure. We have $0 \leq r_p(A) \leq \dim(A)$. The number $r_p(A)$ is invariant under isogenies over k , and satisfies

$r_p(A_1 \times A_2) = r_p(A_1) + r_p(A_2)$. An elliptic curve over k is called ordinary if its p -rank is 1 and is called supersingular if its p -rank is 0. In fact, an elliptic curve is supersingular if and only if $p|a_1$ where $f(t) = x^2 + a_1x + q$ is its characteristic polynomial. An abelian variety A of dimension g over k is called ordinary if its p -rank is g and supersingular if it is $\bar{k}$ -isogenous to a power of a supersingular elliptic curve. A curve C over $\mathbb{F}_q$ is called supersingular if the Jacobian $J_C(\mathbb{F}_q)$ of C is supersingular.

We also will compare our curves to the following famous bound.

Theorem 2.2. (*Hasse-Weil- Serre Bound*) *If C is a curve of genus g over $\mathbb{F}_q$, then*

$$|\#C(\mathbb{F}_q) - (q + 1)| \leq \lfloor 2\sqrt{q} \rfloor$$

The curve C is said to have defect Δ if

$$\#C(\mathbb{F}_q) = q + 1 + \lfloor 2\sqrt{q} \rfloor - \Delta.$$

2.3 Kani-Rosen Decomposition

If $G \leq \text{Aut}(C)$ is a automorphism group of a curve C , then any $H \leq G$ defines an idempotent $\epsilon_H \in \mathbb{Q}[G]$ by

$$\epsilon_H := \frac{1}{|H|} \sum_{h \in H} h.$$

Definition. For $\epsilon_1, \epsilon_2 \in \text{End}_0(J_C) := \text{End}(J_C) \otimes_{\mathbb{Z}} \overline{\mathbb{Q}}$, we say $\epsilon_1 \sim \epsilon_2$ if for all $\overline{\mathbb{Q}}$ -characters χ of $\text{End}_0(J_C)$. we have $\chi(\epsilon_1) = \chi(\epsilon_2)$.

Also $\epsilon_H(J_C) \sim J_{C/H}$.

Theorem 2.3. (Kani-Rosen) *Given a curve C , let $G \leq \text{Aut}(C)$ be a finite group. Given $\epsilon_1, \dots, \epsilon_n, \epsilon'_1, \dots, \epsilon'_m \in \text{End}_0(J_C)$ the idempotent relation*

$$\epsilon_1 + \dots + \epsilon_n \sim \epsilon'_1 + \dots + \epsilon'_m$$

holds in $\text{End}_0(J_C)$ if and only if we have following isogeny relation

$$\epsilon_1(J_C) + \dots + \epsilon_n(J_C) \sim \epsilon'_1(J_C) + \dots + \epsilon'_m(J_C).$$

3 Invariant theory and Quotient Curves

Here we present our algorithm for computing quotient curves of our given curve, using Gröbner bases.

Let $K[X] := K[x_1, \dots, x_n]$ be a polynomial ring in n variables over the field K . Let Γ be a sub-group of the group $GL_n(K)$ of invertible $n \times n$ -matrices with entries from K . Then

$$K[x_1, \dots, x_n]^\Gamma := \{f \in K[x_1, \dots, x_n] \mid f = f \circ \pi \text{ for all } \pi \in \Gamma\}.$$

is a subring of $K[X]$. The Reynolds operator R_Γ of the group Γ is defined as

$$R_\Gamma : K[X] \rightarrow K[X]^\Gamma, \quad R_\Gamma(f) := \frac{1}{|\Gamma|} \sum_{\pi \in \Gamma} f \circ \pi$$

Theorem 3.1. (Hilbert's finiteness theorem) *The invariant ring $(K[X])^\Gamma$ of a finite matrix group $\Gamma \in GL_n(K)$ is finitely generated.*

Theorem 3.2. (Noether's degree bound) *The invariant ring $K[x]^\Gamma$ of finite matrix group Γ has an algebra basis consisting of at most $\binom{n+|\Gamma|}{n}$ invariants whose degree is bounded above by the group order $|\Gamma|$.*

To help find the exact number of number of invariants we have following results

Theorem 3.3. (Molien) *The Hilbert series of the invariant ring of $K[x]^\Gamma$ equals*

$$\Phi_\Gamma(z) = \frac{1}{|\Gamma|} \sum_{\pi \in \Gamma} \frac{1}{\det(id - z\pi)}.$$

The following result may or may not be new; we have not found a reference.

Theorem 3.4. *Let $\text{char}(K) \nmid |\Gamma|$ and $I \subset K[X]$ be ideal fixed by Γ . Then*

$$\frac{K[X]^\Gamma}{I \cap K[X]^\Gamma} \cong \left(\frac{K[X]}{I}\right)^\Gamma.$$

Proof. Define a map

$$\Pi : K[X]^\Gamma \rightarrow \left(\frac{K[X]}{I}\right)^\Gamma, \quad f \mapsto f + I.$$

We claim that this map is surjective. For all $\pi \in \Gamma$ we have $(f + I)^\pi := f^\pi + I$. Let $f + I \in \left(\frac{K[X]}{I}\right)^\Gamma$. This implies $f^\pi + I = f + I$ for all $\pi \in \Gamma$. As I is fixed by Γ and Γ is finite, summing over all the elements of Γ we get $|\Gamma|f + I = \sum_{\pi \in \Gamma} f^\pi + I$. Since $|\Gamma| \nmid \text{char} K$, we have $f + I = R_\Gamma(f) + I$. As $R_\Gamma(f) \in K[X]^\Gamma$ the map defined above is surjective. For kernel, $\Pi(f) = I$ which is possible if and only if $f \in I \cap K[X]^\Gamma$, which proves the assertion. $\square$

Theorem 3.5. *Let $K(V) := \text{Quot}(K[V])$ be a rational function field on vector space V . Then $K(V)^\Gamma = \text{Quot}(K[V]^\Gamma)$.*

Proof. See [14] 3.7.8. □

Definition. Let C be a curve and let Γ be a subgroup of the automorphism group $\text{Aut}(C)$. Then quotient curve C/Γ is the curve whose function field is given by

$$K(C/\Gamma) := \{f \in K(C) \mid f = f \circ \pi \text{ for all } \pi \in \Gamma\}.$$

Here is our algorithm for computing a quotient curve, which we will make use of in this article.

Algorithm 1. *Input:* A curve C and Γ a subgroup of the automorphism group $\text{Aut}(C)$. *Output:* The set of polynomials defining the quotient variety C/Γ .

1. Compute the set of polynomials F invariant under Γ defining the variety C .
2. Compute fundamental set of invariants $\{I_1(x), \dots, I_r(x)\}$.
3. Compute Gröbner basis G_0 for the ideal generated by $\{I_1(x) - y_1, \dots, I_r(x) - y_r\}$ in $K[x_1, \dots, x_n, y_1, \dots, y_r]$.
4. Compute Gröbner basis G_1 of $F \cup G_0$.
5. Compute $H := G_1 \cap K[y_1, \dots, y_r]$. Then $V(H) = C/\Gamma$.

It is clear from the definition of the curve we are studying, which is defined by $x^2 + y^2 + z^2 + w^2 = 0$, $x^3 + y^3 + z^3 + w^3 = 0$, that the symmetric group S_4 is a subgroup of the automorphism group of C . We are going to see later that the following quotient curves are important:

$$\begin{aligned} &C/(1, 2) \\ &C/(1, 2, 3) \\ &C/(1, 2, 3, 4). \end{aligned}$$

Here we identify 1 with x , 2 with y , and so on, so the permutation $(1, 2, 3)$ denotes the map that sends x to y , and y to z , and z to x , and fixes w .

The next sections study these quotients, using Algorithm 1.

4 First Things

For the rest of this paper, C will denote the curve in $\mathbb{P}^3(\overline{\mathbb{Q}})$ defined by the two equations

$$\begin{aligned} F_1(x, y, z, w) &= x^2 + y^2 + z^2 + w^2 = 0 \\ F_2(x, y, z, w) &= x^3 + y^3 + z^3 + w^3 = 0. \end{aligned}$$

Lemma 4.1. *The curve C is nonsingular over $\overline{\mathbb{Q}}$, and C has good reduction modulo p for all $p \geq 5$.*

Proof. It is enough to prove the result for an affine version (say $w = 1$) of C , which is non-singular if and only if

$$\begin{bmatrix} \frac{\partial F_1}{\partial x} & \frac{\partial F_1}{\partial y} & \frac{\partial F_1}{\partial z} \\ \frac{\partial F_2}{\partial x} & \frac{\partial F_2}{\partial y} & \frac{\partial F_2}{\partial z} \end{bmatrix}$$

has full rank. Consider the submatrix

$$\begin{bmatrix} \frac{\partial F_1}{\partial x} & \frac{\partial F_1}{\partial y} \\ \frac{\partial F_2}{\partial x} & \frac{\partial F_2}{\partial y} \end{bmatrix} = \begin{bmatrix} 2x & 2y \\ 3x^2 & 3y^2 \end{bmatrix}$$

with determinant $6xy(x - y) = 0$. If $p \neq 2, 3$ this implies $x = y$ or $x = 0$ or $y = 0$. Similarly considering the determinants of other 2×2 submatrices we get $x = y = z = 0$ which is not a point on the curve. $\square$

Theorem 4.2. *The genus of C is 4.*

Proof. Eliminating z from affine equation of the curve above, we get an irreducible plane curve with some singularities given by

$$f := (x^3 + y^3 + 1)^2 + (x^2 + y^2 + 1)^3 = 0.$$

For singularities, $\frac{\partial f}{\partial x} = 0$, $\frac{\partial f}{\partial y} = 0$ and solving these gives

$$x^3 + y^3 + 1 = 0, x^2 + y^2 + 1 = 0.$$

There are six solutions to these equations, they are the points $(a, (1 + a^3)/(1 + a^2))$ where a is a root of $2x^6 + 3x^4 + 2x^3 + 3x^2 + 2$.

Let $P = (a, b)$ be a singularity. Then $f(x + a, y + b) = ((x + a)^3 + (y + b)^3 + 1)^2 + ((x + a)^2 + (y + b)^2 + 1)^3 = F_0 + F_1 + \dots + F_6$. where F_i is a form of degree i .

We have $F_0 = F_1 = 0$ and $F_2 = (3a^2x + 3b^2y)^2 \neq 0$ so the multiplicity of f at P is 2. Since P was arbitrary $v_{P_i} = \text{Mult}_{P_i} C = 2$. By the genus formula [16], 148,

$$\text{genus}(C) = \frac{(n-1)(n-2)}{2} - \sum_{i=1}^6 \frac{(v_{P_i})(v_{P_i} - 1)}{2} = 10 - 6 = 4.$$

$\square$

We note for the record that the curve C is absolutely irreducible over $\mathbb{Q}$ and $\mathbb{F}_p$ for $p \geq 5$, but we will not use it in this paper.

5 Genus 1 and Genus 0 Quotients

We discuss the affine model of C with $w = 1$.

Theorem 5.1. *The quotient curve $C/(1, 2)$ is an elliptic curve over $\overline{\mathbb{Q}}$.*

Proof. We have $(1, 2) := \begin{cases} x \mapsto y \\ y \mapsto x. \end{cases}$

This can be done directly but we proceed to use Algorithm 1. The fundamental invariants are $a := x + y$, $b := z$, $c := x^2 + y^2$. The Groebner basis is found with MAGMA to be $G_0 = \{x + y - a, y^2 - ya + \frac{1}{2}a^2 - \frac{1}{2}c, z - b\}$ and $G_1 := \{x + y - a, y^2 - ya + \frac{1}{2}a^2 - \frac{1}{2}c, z - b, a^3 - 3ac + 2bc + 2b - 2, b^2 + c + 1\}$. which gives $G_2 = \{a^3 - 3ac + 2bc + 2b - 2, b^2 + c + 1\}$. Here G_2 is the defining ideal for the $K(C/(1, 2))$. Substituting the second equation of G_2 into the first gives $a^3 + 3ab^2 + 3a - 2b^3 - 2 = 0$ which on homegenization gives $a^3 + 3ab^2 + 3aw^2 - 2b^3 - 2w^3 = 0$. Under the bi-rational substitution $x = 3a$, $y = 9b$, $z = -a + b + w$ and putting $z = 1$ we get $y^2 - 3xy - 9y = x^3 - \frac{27}{2}x - 27$. $\square$

Theorem 5.2. *The quotient curve $C/(1, 2, 3, 4)$ is an elliptic curve.*

Proof. This time the Gröbner basis is too large to include. MAGMA gives $C/(1, 2, 3, 4)$ as an elliptic curve defined by

$$y^2 - 96xy + 110592y = x^3 + 3456x^2 + 14598144x - 5718933504$$

over $\overline{\mathbb{Q}}$. $\square$

Theorem 5.3. *The quotient curves $C/(1, 2)$ and $C/(1, 2, 3, 4)$ are isomorphic over $\overline{\mathbb{Q}}$.*

Proof. Both the quotient curves have the same j -invariant equal to -36. Hence they are isomorphic over $\overline{\mathbb{Q}}$. In fact, the isomorphism is defined over $\mathbb{Q}$ and is given by

$$\phi : C/(1, 2) \rightarrow C/(1, 2, 3, 4), \quad (x, y) \rightarrow (1024x - 1152, 32768y - 2580481).$$

Their Weierstrass form is given by

$$y^2 = x^3 - 27x - 378.$$

$\square$

Remark: MAGMA tells us that the endomorphism ring of $C/(1, 2)$ is $\mathbb{Z}$, and so this elliptic curve does not have complex multiplication.

Theorem 5.4. *The genus of the quotient curve C/S_4 is 0.*

Proof. The fundamental invariants are $a := x + y + z + w$, $b := x^2 + y^2 + z^2 + w^2$, $c := x^3 + y^3 + z^3 + w^3$, $d := x^4 + y^4 + z^4 + w^4$.

The Groebner Basis $G_0 = \{x + y + z + w - a, y^2 + yz + yw - ya + z^2 + zw - za + w^2 - wa + 1/2a^2 - \frac{1}{2}b, z^3 + z^2w - z^2a + zw^2 - zwa + \frac{1}{2}za^2 - \frac{1}{2}zb + w^3 - w^2a + 1/2wa^2 - \frac{1}{2}wb - \frac{1}{6}a^3 + \frac{1}{2}ab - \frac{1}{3}c, w^4 - w^3a + 1/2w^2a^2 - \frac{1}{2}w^2b - \frac{1}{6}wa^3 + \frac{1}{2}wab - \frac{1}{3}wc + \frac{1}{24}a^4 - \frac{1}{4}a^2b + \frac{1}{3}ac + \frac{1}{8}b^2 - \frac{1}{4}d\}$.
 $G_1 = \{x + y + z + w - a, y^2 + yz + yw - ya + z^2 + zw - za + w^2 - wa + \frac{1}{2}a^2, z^3 + z^2w - z^2a + zw^2 - zwa + \frac{1}{2}za^2 + w^3 - w^2a + \frac{1}{2}wa^2 - \frac{1}{6}a^3, w^4 - w^3a + \frac{1}{2}w^2a^2 - \frac{1}{6}wa^3 + \frac{1}{24}a^4 - \frac{1}{4}d, b, c\}$,
 which gives $G_2 = \{b, c\}$. The equations from G_2 define a projective line as asserted. $\square$

6 Genus 2 Quotient

Theorem 6.1. *The quotient curve $C/(1, 2, 3)$ is a hyperelliptic curve of genus 2.*

Proof. We have $(1, 2, 3) := \begin{cases} x \mapsto y \\ y \mapsto z \\ z \mapsto x. \end{cases}$

The fundamental invariants are $a := x + y + z$, $b := x^2 + y^2 + z^2$, $c := x^3 + y^3 + z^3$, $d := x^2z + xy^2 + yz^2$.

MAGMA gives the Groebner Basis G_1 to be $\{x + y + z - a, y^2 - \frac{4}{3}yd^5 + \frac{16}{3}yd^4 - \frac{28}{3}yd^3 + \frac{28}{3}yd^2 - \frac{17}{6}yd - \frac{1}{6}y + \frac{1}{24}z^2a^5d - \frac{1}{12}z^2a^5 - \frac{1}{12}z^2a^4d^2 + \frac{5}{24}z^2a^4d - \frac{1}{12}z^2a^4 + \frac{1}{6}z^2a^3d^3 - \frac{1}{2}z^2a^3d^2 + \frac{3}{4}z^2a^3d - \frac{5}{6}z^2a^3 - \frac{1}{3}z^2a^2d^4 + \frac{7}{6}z^2a^2d^3 - 2z^2a^2d^2 + \frac{25}{12}z^2a^2d - \frac{1}{6}z^2a^2 + \frac{1}{2}z^2ad^3 - \frac{3}{2}z^2ad^2 + \frac{17}{8}z^2ad - \frac{5}{2}z^2a - z^2d^4 + \frac{7}{2}z^2d^3 - \frac{19}{4}z^2d^2 + \frac{25}{8}z^2d + \frac{3}{2}z^2 - \frac{1}{48}za^5 + \frac{1}{24}za^4d - \frac{1}{48}za^4 - \frac{1}{12}za^3d^2 + \frac{1}{12}za^3d + \frac{1}{24}za^3 + \frac{1}{6}za^2d^3 - \frac{1}{4}za^2d^2 + \frac{5}{24}za^2 - \frac{1}{3}zad^4 + \frac{2}{3}zad^3 - \frac{1}{4}zad^2 - \frac{5}{12}zad + \frac{55}{48}za - \frac{2}{3}zd^5 + \frac{11}{3}zd^4 - \frac{49}{6}zd^3 + \frac{119}{12}zd^2 - \frac{145}{24}zd - \frac{1}{48}z - \frac{1}{48}a^5 + \frac{1}{24}a^4d - \frac{5}{48}a^4 - \frac{1}{12}a^3d^2 + \frac{1}{4}a^3d - \frac{7}{24}a^3 + \frac{1}{6}a^2d^3 - \frac{7}{12}a^2d^2 + \frac{5}{6}a^2d - \frac{5}{8}a^2 + \frac{2}{3}ad^5 - 3ad^4 + 6ad^3 - \frac{83}{12}ad^2 + \frac{7}{2}ad - \frac{7}{16}a - \frac{1}{3}d^4 + \frac{7}{6}d^3 - \frac{7}{4}d^2 + \frac{29}{24}d + \frac{1}{48}, yz + \frac{4}{3}yd^5 - \frac{16}{3}yd^4 + \frac{28}{3}yd^3 - \frac{28}{3}yd^2 + \frac{29}{6}yd - \frac{5}{6}y - \frac{1}{24}z^2a^5d + \frac{1}{12}z^2a^5 + \frac{1}{12}z^2a^4d^2 - \frac{5}{24}z^2a^4d + \frac{1}{12}z^2a^4 - \frac{1}{6}z^2a^3d^3 + \frac{1}{2}z^2a^3d^2 - \frac{3}{4}z^2a^3d + \frac{5}{6}z^2a^3 + \frac{1}{3}z^2a^2d^4 - \frac{7}{6}z^2a^2d^3 + 2z^2a^2d^2 - \frac{25}{12}z^2a^2d + \frac{2}{3}z^2a^2 - \frac{1}{2}z^2ad^3 + \frac{3}{2}z^2ad^2 - \frac{17}{8}z^2ad + \frac{5}{2}z^2a + z^2d^4 - \frac{7}{2}z^2d^3 + \frac{19}{4}z^2d^2 - \frac{25}{8}z^2d + z^2 + \frac{1}{48}za^5 - \frac{1}{24}za^4d + \frac{1}{48}za^4 + \frac{1}{12}za^3d^2 - \frac{1}{12}za^3d - \frac{1}{24}za^3 - \frac{1}{6}za^2d^3 + \frac{1}{4}za^2d^2 - \frac{5}{24}za^2 + \frac{1}{3}zad^4 - \frac{2}{3}zad^3 + \frac{1}{4}zad^2 + \frac{5}{12}zad - \frac{55}{48}za + \frac{2}{3}zd^5 - \frac{11}{3}zd^4 + \frac{49}{6}zd^3 - \frac{119}{12}zd^2 + \frac{169}{24}zd - \frac{95}{48}z + \frac{1}{48}a^5 - \frac{1}{24}a^4d + \frac{5}{48}a^4 + \frac{1}{12}a^3d^2 - \frac{1}{4}a^3d + \frac{7}{24}a^3 - \frac{1}{6}a^2d^3 + \frac{7}{12}a^2d^2 - \frac{5}{6}a^2d + \frac{5}{8}a^2 - \frac{2}{3}ad^5 + 3ad^4 - 6ad^3 + \frac{83}{12}ad^2 - \frac{9}{2}ad + \frac{23}{16}a + \frac{1}{3}d^4 - \frac{7}{6}d^3 + \frac{7}{4}d^2 - \frac{29}{24}d + \frac{17}{48}, ya + 2yd - y + \frac{1}{2}z^2a^2 + \frac{3}{2}z^2 + za + zd - 2z - \frac{1}{2}a^2 - ad + a + \frac{1}{2}, yd^6 - 3yd^5 + 6yd^4 - 6yd^3 + \frac{33}{32}yd^2 - \frac{3}{4}yd + \frac{1}{8}y - \frac{1}{32}z^2a^5d^2 + \frac{1}{32}z^2a^5d - \frac{1}{32}z^2a^5 + \frac{1}{16}z^2a^4d^3 - \frac{3}{32}z^2a^4d^2 + \frac{3}{32}z^2a^4d - \frac{1}{32}z^2a^4 - \frac{1}{8}z^2a^3d^4 + \frac{1}{4}z^2a^3d^3 - \frac{9}{16}z^2a^3d^2 + \frac{7}{16}z^2a^3d - \frac{5}{16}z^2a^3 + \frac{1}{4}z^2a^2d^5 - \frac{5}{8}z^2a^2d^4 + \frac{11}{8}z^2a^2d^3 - \frac{19}{16}z^2a^2d^2 + \frac{13}{16}z^2a^2d - \frac{1}{16}z^2a^2 - \frac{3}{8}z^2ad^4 + \frac{3}{4}z^2ad^3 - \frac{51}{32}z^2ad^2 + 4532z^2ad - \frac{27}{32}z^2a + \frac{3}{4}z^2d^5 - \frac{15}{8}z^2d^4 + \frac{51}{16}z^2d^3 - \frac{69}{32}z^2d^2 + \frac{27}{32}z^2d + \frac{21}{32}z^2 + \frac{1}{64}za^5d + \frac{1}{64}za^5 - \frac{1}{32}za^4d^2 - \frac{1}{64}za^4d + \frac{1}{64}za^4 + \frac{1}{16}za^3d^3 + \frac{3}{32}za^3d + \frac{5}{32}za^3 - \frac{1}{8}za^2d^4 + \frac{1}{16}za^2d^3 - \frac{3}{16}za^2d^2 - \frac{11}{32}za^2d + \frac{1}{32}za^2 + \frac{1}{4}zad^5 - \frac{1}{4}zad^4 + \frac{7}{16}zad^3 + \frac{1}{2}zad^2 + \frac{1}{64}zad + \frac{29}{64}za + \frac{1}{2}zd^6 - \frac{9}{4}zd^5 + \frac{39}{8}zd^4 - \frac{105}{16}zd^3 + \frac{159}{32}zd^2 - \frac{129}{64}zd - \frac{11}{64}z + \frac{1}{64}a^5d - \frac{1}{64}a^5 - \frac{1}{32}a^4d^2 + \frac{3}{64}a^4d - \frac{1}{64}a^4 + \frac{1}{16}a^3d^3 - \frac{1}{8}a^3d^2 + \frac{7}{32}a^3d - \frac{5}{32}a^3 - \frac{1}{8}a^2d^4 + \frac{5}{16}a^2d^3 - \frac{9}{16}a^2d^2 + \frac{13}{32}a^2d - \frac{1}{32}a^2 - \frac{1}{2}6 +$

$$\frac{7}{4}ad^5 - \frac{15}{4}ad^4 + 7\frac{1}{16}ad^3 - \frac{55}{16}ad^2 + \frac{81}{64}ad - \frac{33}{64}a + \frac{1}{4}d^5 - \frac{5}{8}d^4 + \frac{19}{16}d^3 - \frac{23}{32}d^2 + \frac{19}{64}d + \frac{23}{64}, z^3 - z^2a + \frac{1}{2}za^2 + \frac{1}{2}z - \frac{1}{6}a^3 - \frac{1}{2}a + \frac{1}{3}, a^6 + 9a^4 - 8a^3 + 27a^2 + 24ad - 48a + 24d^2 - 24d + 27, b + 1, c + 1\}.$$

Therefore by Algorithm 1 the defining set for the quotient curve is $G_2 = \{a^6 + 9a^4 - 8a^3 + 27a^2 + 24ad - 48a + 24d^2 - 24d + 27, b + 1, c + 1\}$. The change of variables $y := -\frac{1}{2}a - \frac{3}{2}d$, $x = \frac{1}{2}(-a + 1)$ defines a hyperelliptic curve given by $y^2 + (x^3 + x^2)y = -x^6 + 4x^5 - 25x^4 + 36x^3 - 36x^2 + 18x - 6$.

□

Let V be a curve of genus 2 over a field k . Let $Jac(V)$ be its Jacobian. We say that J is split over k if J is isogenous over k to a product of elliptic curves $E_1 \times E_2$.

Theorem 6.2. (Kuhn[20]) *Let J be a Jacobian of a curve V of genus 2 over a field k of characteristic different from 2. Suppose that J is split. Then there are elliptic curves E_1, E_2 over k , and an integer $n > 1$ such that $E_1[n]$ and $E_2[n]$ are isomorphic as group schemes and J is isogenous to $E_1 \times E_2$. Furthermore, the curve V admits degree n covers $V \rightarrow E_1$ and $V \rightarrow E_2$.*

Whenever the above happens, we call $Jac(V)$ a (n, n) -split, and we say $Jac(V)$ is (n, n) -isogenous to $E_1 \times E_2$.

Let I_2, I_4, I_6 , and I_{10} (see [22]) of a genus 2 curve V . We define absolute invariants,

$$i_1 = 144 \frac{I_4}{I_2^2}, \quad i_2 = -1728 \frac{I_2 I_4 - 3I_6}{I_2^3}, \quad i_3 = 486 \frac{I_{10}}{I_2^5}.$$

Theorem 6.3. (Bruin-Doerksen) *The absolute invariants i_1, i_2, i_3 of a genus 2 curve with optimally $(4, 4)$ -split Jacobian satisfy an equation $L_4 = 0$.*

The equation L_4 is too big to be reproduced here. See [1, 2] for proof and the equation L_4 .

The projective equation of $C/(1, 2, 3)$ is given by $y^2 z^4 + (x^3 z^2 + x^2 z^3)y = -x^6 + 4x^5 z - 25x^4 z^2 + 36x^3 z^3 - 36x^2 z^4 + 18x z^5 - 6z^6$. Performing the birational substitution $(x, y, z) \rightarrow (x, x^3, x^2 z + 2y)$ and then putting $z = 1$ gives the Weierstrass form

$$y^2 = -3x^6 - 18x^5 - 99x^4 - 144x^3 - 144x^2 - 72x - 24.$$

The Igusa invariants are $I_2 = -138240$, $I_4 = 234150912$, $I_6 = -448888946688$, $I_{10} = -12999674453557248$. This gives the absolute invariants to be $i_1 = 2823/1600$, $i_2 = 2597331/128000$, $i_3 = 6561/52428800000$. It can be checked using any computer algebra package that the absolute invariants satisfy the equation $L_4 = 0$. Hence by theorem 6.3 we strongly suspected that $Jac(C/(1, 2, 3))$ is a $(4, 4)$ -split. We now prove this.

Theorem 6.4. *Let E_1 be the elliptic curve $C/(1, 2)$, and let E_2 be the elliptic curve over $\overline{\mathbb{Q}}$ defined by $y^2 + xy = x^3 - x^2 - 6x + 8$. Then Jacobian of $C/(1, 2, 3)$ is $(4, 4)$ -isogenous to $E_1 \times E_2$.*

Proof. We will use the method given in section 4 of [1]. We first prove the existence of $(2, 2)$ Richelot isogeny from $Jac(C/(1, 2, 3))$ to a certain $(2, 2)$ -split Jacobian, say B , and then construct it. The hyperelliptic curve $C/(1, 2, 3)$ is given by $y^2 = -3F$ where $F = x^6 + 6x^5 + 33x^4 + 48x^3 + 48x^2 + 24x + 8$. The splitting field of the polynomial F has is the cubic number field S defined by $x^3 - 3x + 4$ and discriminant $d = -324$. Therefore we get $F = F_1 F_2 F_3$ where

$$\begin{aligned} F_1 &= x^2 + \frac{1}{18}(a^4 - 3a^2)x + \frac{1}{18}(a^4 - 9a^2 - 18), \\ F_2 &= x^2 + \frac{1}{36}(-a^5 - a^4 + 15a^3 + 3a^2 - 72a + 108)x + \frac{1}{72}(-a^5 - 2a^4 + 15a^3 + 18a^2 - 108a - 72), \\ F_3 &= x^2 + \frac{1}{36}(a^5 - a^4 - 15a^3 + 3a^2 + 72a + 108)x + \frac{1}{72}(a^5 - 2a^4 - 15a^3 + 18a^2 + 108a - 72), \end{aligned}$$

where a satisfies $x^6 - 18x^4 + 81x^2 + 324$. If we write $F_j(X) = q_2X^2 + q_{1,j}X + q_{0,j}$ then we define

$$\delta := \det \begin{pmatrix} q_{0,1} & q_{1,1} & q_2 \\ q_{0,2} & q_{1,2} & q_2 \\ q_{0,3} & q_{1,3} & q_2 \end{pmatrix} = 2a^3 - 18a.$$

Since $\delta \neq 0$, B is the Jacobian of genus 2 curve. We will find the equation of a curve $\tilde{C}$ such that $Jac(\tilde{C}) = B$. We define for $(i, j, k) = (1, 2, 3), (2, 3, 1), (3, 1, 2)$

$$G_i(X) = \delta^{-1} \det \begin{pmatrix} \frac{d}{dX}F_j(X) & \frac{d}{dX}F_k(X) \\ F_j(X) & F_k(X) \end{pmatrix}.$$

Hence, $G_1 = \frac{1}{648}(-a^4 + 27a^2 - 108)x^2 + \frac{1}{216}(-a^4 + 15a^2 - 36)x + \frac{1}{162}(-a^4 + 18a^2 - 108)$,
 $G_2 = \frac{1}{1296}(-a^5 + a^4 + 15a^3 - 27a^2 + 108)x^2 + \frac{1}{432}(a^4 - 15a^2 + 36a + 36)x + \frac{1}{1296}(-a^5 + 4a^4 + 15a^3 - 72a^2 + 108a - 216)$,
and $G_3 = \frac{1}{1296}(a^5 + a^4 - 15a^3 - 27a^2 + 108)x^2 + \frac{1}{432}(a^4 - 15a^2 - 36a + 36)x + \frac{1}{1296}(a^5 + 4a^4 - 15a^3 - 72a^2 - 108a - 216)$.

Hence we get $\tilde{C}$ defined by $y^2 = dG_1G_2G_3$, i.e., $y^2 = f(x) = 2x^6 + 6x^5 + 15x^4 + 18x^3 + 15x^2 + 6x + 2$.

To see that $\tilde{C}$ is a $(2, 2)$ cover of elliptic curves (say E_1 and E_2) make the change of variable $x \rightarrow \frac{x+1}{x-1}$ we get $f = \frac{64x^6+36x^4+24x^2+4}{x^6-6x^5+15x^4-20x^3+15x^2-6x+1}$. Put $u = x^2$ in the numerator of f we get $y^2 = 64u^3 + 36u^2 + 24u + 4$ as one of the elliptic curve say E_1 , the covering map is given by $(x, y) \rightarrow (x^2, y)$. Then by [7] p.155, E_2 is given by $y^2 = 4u^3 + 24u^2 + 36u + 64$ and the covering map is given by $(x, y) \rightarrow (\frac{1}{x^2}, \frac{y}{x^3})$. □

7 Arithmetic Properties of The Quotients that arise from the Splitting

When we write $L(J_C)$ we mean the L-polynomial of the reduction modulo p of the Jacobian of C . The Jacobian has good reduction because C does.

Theorem 7.1. *Over $\mathbb{F}_p$, $p \geq 5$,*

$$L(J_C) = L(J_{C/(1,2)}) \times L(J_{C/(1,2,3)}) \times L(J_{C/(1,2,3,4)}).$$

Proof. First we will work over $\overline{\mathbb{Q}}$. By Lemma 4.2, C is a non-singular over $\overline{\mathbb{Q}}$. The automorphism group of C contains the symmetric group on 4 symbols i.e; $S_4 \leq \text{Aut}(C)$. By the symmetry of the curve, one can conclude that $\epsilon_{\langle\sigma\rangle}(J_C) \sim \epsilon_{\langle\tau\rangle}(J_C)$ if σ and τ are of same cycle type.

In S_4 there are

1. six cyclic subgroups of order 2, say $\sigma_1, \dots, \sigma_6$.
2. four cyclic subgroups of order 3, say $\tau_1, \dots, \tau_4$.
3. three cyclic subgroups of order 4, say $F_1, \dots, F_3$.

We have the following idempotent relation in S_4 , namely

$$12\epsilon_{id} + 24\epsilon_{S_4} = 2[\epsilon_{\sigma_1} + \dots \epsilon_{\sigma_6}] + 3[\epsilon_{\tau_1} + \dots \epsilon_{\tau_4}] + 4[\epsilon_{F_1} + \dots \epsilon_{F_3}],$$

which by Kani-Rosen decomposition implies

$$\epsilon_{id}(J_C)^{12} \times \epsilon_{S_4}(J_C)^{24} \sim \epsilon_{\sigma_1}(J_C)^2 \times \dots \times \epsilon_{\sigma_6}(J_C)^2 \times \epsilon_{\tau_1}(J_C)^3 \times \dots \times \epsilon_{\tau_4}(J_C)^3 \times \epsilon_{F_1}(J_C)^4 \times \dots \times \epsilon_{F_3}(J_C)^4.$$

This gives the following isogeny over $\overline{\mathbb{Q}}$

$$J_C^{12} \times J_{C/S_4}^{24} \sim J_{C/(1,2)}^{12} \times J_{C/(1,2,3)}^{12} \times J_{C/(1,2,3,4)}^{12}.$$

Reducing this isogeny modulo $p \geq 5$ gives an isogeny over $\mathbb{F}_p$, which means that the L -polynomials satisfy

$$L(J_C)^{12} \times L(J_{C/S_4})^{12} = L(J_{C/(1,2)})^{12} \times L(J_{C/(1,2,3)})^{12} \times L(J_{C/(1,2,3,4)})^{12}.$$

But $\dim(J_{C/S_4}) = 0$ from Theorem 5.4 which implies that $L((J_{C/S_4})) = 1$. By taking the 12th-root on both sides we get the result. $\square$

Corollary 7.2. *The modulo p reductions of J_C and $J_{C/(1,2)} \times J_{C/(1,2,3)} \times J_{C/(1,2,3,4)}$ are isogenous over $\mathbb{F}_p$.*

Proof. By Theorem 7.1 and Tate's theorem, we can conclude that there is an $\mathbb{F}_p$ -isogeny between (the mod p reductions of) J_C and $J_{C/(1,2)} \times J_{C/(1,2,3)} \times J_{C/(1,2,3,4)}$. $\square$

From the equations computed earlier of the quotient curves defined over $\overline{\mathbb{Q}}$, we tabulate their L -polynomials over $\mathbb{F}_p$ for few primes p below.

p	$L(J_{C/(1,2)})$	$L(J_{C/(1,2,3)})$	$L(J_{C/(1,2,3,4)})$	p -rank
5	$(5t^2 - t + 1)$	$(5t^2 - t + 1)(5t^2 + 3t + 1)$	$(5t^2 - t + 1)$	4
7	$7t^2 + 1$	$(7t^2 + 1)(7t^2 + 4t + 1)$	$7t^2 + 1$	1
11	$11t^2 - 4t + 1$	$(11t^2 - 4t + 1)(11t^2 + 1)$	$11t^2 - 4t + 1$	3
13	$13t^2 + 5t + 1$	$(13t^2 + t + 1)(13t^2 + 5t + 1)$	$13t^2 + 5t + 1$	4
17	$17t^2 - 5t + 1$	$(17t^2 - 5t + 1)(17t^2 + 3t + 1)$	$17t^2 - 5t + 1$	4
19	$19t^2 - 8t + 1$	$(19t^2 - 8t + 1)(19t^2 + 4t + 1)$	$19t^2 - 8t + 1$	4
23	$23t^2 - 4t + 1$	$(23t^2 - 4t + 1)(23t^2 + 1)$	$23t^2 - 4t + 1$	3
29	$29t^2 + 3t + 1$	$(29t^2 - 9t + 1)(29t^2 + 3t + 1)$	$29t^2 + 3t + 1$	4
31	$31t^2 + 4t + 1$	$(31t^2 + 4t + 1)^2$	$31t^2 + 4t + 1$	4
37	$37t^2 - 3t + 1$	$(37t^2 + t + 1)(37t^2 - 3t + 1)$	$37t^2 - 3t + 1$	4
37	$37t^2 - 3t + 1$	$(37t^2 - 3t + 1)(37t^2 + t + 1)$	$37t^2 - 3t + 1$	4
41	$41t^2 - 6t + 1$	$(41t^2 - 6t + 1)^2$	$41t^2 - 6t + 1$	4
43	$43t^2 - 4t + 1$	$(43t^2 - 8t + 1)(43t^2 - 4t + 1)$	$43t^2 - 4t + 1$	4
47	$47t^2 - 12t + 1$	$(47t^2 - 12t + 1)(47t^2 + 12t + 1)$	$47t^2 - 12t + 1$	4
53	$53t^2 - 10t + 1$	$(53t^2 - 10t + 1)(53t^2 + 6t + 1)$	$53t^2 - 10t + 1$	4
59	$59t^2 + 8t + 1$	$(59t^2 + 1)(59t^2 + 8t + 1)$	$59t^2 + 8t + 1$	3
61	$61t^2 + 5t + 1$	$(61t^2 + t + 1)(61t^2 + 5t + 1)$	$61t^2 + 5t + 1$	4
67	$67t^2 - 8t + 1$	$(67t^2 - 8t + 1)(67t^2 + 4t + 1)$	$67t^2 - 8t + 1$	4
71	$71t^2 + 16t + 1$	$(71t^2 + 12t + 1)(71t^2 + 16t + 1)$	$71t^2 + 16t + 1$	4
73	$73t^2 + 5t + 1$	$(73t^2 - 11t + 1)(73t^2 + 5t + 1)$	$73t^2 + 5t + 1$	4
79	$79t^2 - 4t + 1$	$(79t^2 - 4t + 1)(79t^2 + 16t + 1)$	$79t^2 - 4t + 1$	4
83	$83t^2 + 4t + 1$	$(83t^2 + 4t + 1)(83t^2 + 12t + 1)$	$83t^2 + 4t + 1$	4
89	$89t^2 + 3t + 1$	$(89t^2 + 3t + 1)^2$	$89t^2 + 3t + 1$	4
97	$97t^2 - 2t + 1$	$(97t^2 - 2t + 1)^2$	$97t^2 - 2t + 1$	4
101	$101t^2 + 6t + 1$	$(101t^2 + 6t + 1)^2$	$101t^2 + 6t + 1$	4
103	$103t^2 - 4t + 1$	$(103t^2 - 4t + 1)(103t^2 + 4t + 1)$	$103t^2 - 4t + 1$	4

Remark: Theorem 6.4 implies that over $\mathbb{F}_p$, for all $p \geq 5$, the Jacobian of $C/(1, 2, 3)$ (reduced modulo p) is isogenous to the product of $\overline{E_1}$ and $\overline{E_2}$, where $\overline{E}$ denotes the reduction of E modulo p . By Corollary 7.2 we get that

$$\overline{J_C} \sim \overline{E_1}^3 \times \overline{E_2}.$$

Remark: For $p = 31, 41, 89, 97, 101$, the table shows that $\overline{E_1}$ is isogenous to $\overline{E_2}$, and so

$$\overline{J_C} \sim \overline{E_1}^4.$$

It would be interesting to know how often this happens. If $\overline{J_C} \sim \overline{E_1} \times \overline{E_2}$ has L-polynomial $q^2t^4 + qa_1t^3 + a_2t^2 + a_1t + 1$ then $\overline{E_1}$ is isogenous to $\overline{E_2}$ if and only if $a_1^2 - 4a_2 + 8q = 0$, but we cannot see how to predict for which p this will happen.

8 Number of Rational Points

It would be nice to have a formula for the number of rational points on the reduction of C modulo p . However, our investigations show that this is unlikely.

The table below gives the number of rational points of C for some values of q and the Hasse -Weil-Serre bounds.

p	Lower bound	$\#C(\mathbb{F}_q)$	Upper bound
5	-10	6	22
7	-12	12	28
11	-12	0	36
13	-14	30	42
17	-14	6	50
19	-12	0	52
23	-12	12	60
29	-10	30	70
31	-12	48	76
37	-10	30	86
41	-6	18	90
43	-8	24	96
47	-4	24	100
53	-2	30	110
59	0	84	120
61	2	78	122
67	4	48	132
71	8	132	136
73	6	78	142
79	12	84	148
83	12	108	156
89	18	102	162
97	22	90	174
101	22	126	182
103	24	96	184

Remark: 1. For $p = 71$ the defect is only 4, which is very good. We do not know if this is best possible for genus 4 over that field.

2. For $p = 11, 19$ the curve C has no points.

Because we have taken explicit exponents $a = 2, b = 3$ in this article, many things can be computed explicitly. In future work we will investigate other values of a and b .

References

- [1] Nils Bruin, Kevin Doerksen, The arithmetic of genus two curves with (4,4)-split Jacobians, preprint arXiv:0902.3480, 2009.
- [2] Nils Bruin and Kevin Doerksen, *Electronic resources*, <http://www.cecm.sfu.ca/~nbruin/splitigusa>.
- [3] N. Boston, G. McGuire, The Weight Distributions of Cyclic Codes with Two Zeros and Zeta Functions, *Journal of Symbolic Computation* Vol 45, Issue 7, July 2010, 723–733.
- [4] J. Paulhus, Decomposing Jacobians of curves with extra automorphisms. *Acta Arith.* 132 (2008), no. 3, 231-244.
- [5] S. Cutkosky, Resolution of Singularities, AMS Bookstore, 2004.
- [6] Gary McGuire, On certain 3-weight cyclic codes having symmetric weights and a conjecture of Helleseth, Proc. SETA'01, DMTCS Series, Springer, (2001)281-295.
- [7] J. W. S. Cassels and E. V. Flynn, *Prolegomena to a middlebrow arithmetic of curves of genus 2*, London Mathematical Society Lecture Note Series, vol. 230, Cambridge University Press, Cambridge, 1996.
- [8] Iwan Duursma, From Weight Enumerators to Zeta Functions, DAMATH, (1999)
- [9] H. Janwa, G. McGuire and R.M. Wilson, Double-error-correcting cyclic codes and absolutely irreducible polynomials over $GF(2)$, *J. Alg.* **178** (1995) 665–676.
- [10] Wieb Bosma, John Cannon and Catherine Playoust. The Magma algebra system. I. The user language. *J. Symbolic Comput.*, 24(3-4):235-265, 1997.
- [11] J. Wolfmann, The number of solutions of certain diagonal equations over finite fields. *J. Number Theory* 42 (1992), pp. 247-257.
- [12] M.P.Knapp, System of diagonal equations over p-adic fields. *Journal of the London Mathematical Society*, 63(2001) , pp 257-267
- [13] E.Kani and M.Rosen. Idempotent relations and factors of Jacobians, *Math. Ann.*, 284(2):307-327, 1989.
- [14] H.Derksen and G.Kemper. Computational Invariant Theory, *Encyclopedia of Mathematical Sciences*, Vol. 130, Springer 2002.
- [15] D.J.Benson. Polynomial Invariant Of Finite Groups, Cambridge University Press, 1993.

- [16] Shreeram Abhyankar, Algebraic Geometry for Scientists and Engineers, Mathematical Surveys and Monographs, Vol.35,AMS, 1980.
- [17] T.Kasami, The Weight eumerators for several classes of subcodes of second order binary Reed Muller codes, Information and control, 18(1971), 369-394.
- [18] H. Stichtenoth and C. Xing, On the structure of the divisor class group of a class of curves over Finite Fields, Arch. Math., Vol. 65 (1995) 141 -150.
- [19] H.-G. Ruck. Abelian surfaces and Jacobian varieties over finite fields. Compositio Math., 76(3):351-366, 1990.
- [20] Robert M. Kuhn, Curves of genus 2 with split Jacobian, Trans. Amer. Math. Soc. 307 (1988), no.1, 41–49.
- [21] C.Adimoolam, A note on good reduction of simple abelian varieties, Proceedings of the American Mathematical Society, Vol.64, No 2, 196-198.
- [22] Jun-ichi Igusa, Arithmetic variety of moduli for genus two, Ann. of Math. (1960), 612–649.
- [23] John Tate, Endomorphisms of abelian varieties over finite fields, Invent.Math., 2:134-144, 1966.